



Amit Momaya, MD

Orthopedic Surgeon, Chief of Sports Medicine
Team Physician for UAB, Legion FC, and Bulls Hockey
Highlands Appointments: 205.930.8339
MomayaMD.com

PATIENT NAME:

PROCEDURE:

DATE OF SURGERY:

PHYSICAL THERAPY ORDER – MPFL recon with allograft, Tibial Tubercle Osteotomy (AMZ)

Phase I: 0-2 weeks

- Edema and pain control
- Weight bearing status: Heel touch only in extension
- ROM: 0-30 degrees passively
- Brace: locked in extension during ambulation and sleeping
- Exercises: quad sets, gentle patella mobs, calf pumps, SLR

Phase II: 2-6 weeks

- Weight Bearing Status: Heel touch only in extension
- ROM:
 - o 2-4 weeks: 0-60 degrees passively
 - o 4-6 weeks: 0-90 degrees passively
- Brace: per ROM above
- Exercises: Phase I + hip/glutes/core

Phase III: 6-12 weeks

- WB status: slowly progress to full
- ROM: progress to full
- Brace: lateral padded patella brace
- Exercise: Phase II + close chain quads, hamstring curls, step-ups, front/side planks; progressive proprioception exercises

Phase IV: 3-6 months

- WB status: full
- ROM: full
- Brace: lateral padded sleeve as needed
- Exercise: Phase III + advance to open chain exercises; ok for stationary bike at 3-4 mo

Phase V: 6-12 mo

- Brace: none
- Gradual return to sports once cleared by MD