



Amit Momaya, MD

Orthopedic Surgeon, Chief of Sports Medicine Team Physician for UAB, Legion FC, and Bulls Hockey Highlands Appointments: 205.930.8339 MomayaMD.com

PATIENT NAME:

PROCEDURE:

DATE OF SURGERY:

PHYSICAL THERAPY ORDER - MPFL Reconstruction with Allograft

Phase I: 0-2 weeks

- Edema and pain control

- Weight bearing status: partial weight bearing with crutches

- ROM: 0-30 degrees

Brace: locked in extension during ambulation and sleeping

- Exercises: heel slides, gastroc/soleus stretch, calf pumps, straight leg raises

Phase II: 2-6 weeks

Weight Bearing Status:

2-4 weeks: partial WB with brace4-6 weeks: progress to full with brace

- ROM:

2-4 weeks: 0-60 degrees4-6 weeks: 0-90 degrees

- Brace: per ROM above

- Exercises: Phase I + hip/glutes/core

Phase III: 6-12 weeks
- WB status: full

- ROM: full

- Brace: lateral padded patella brace

- Exercise: Phase II + close chain quads, hamstring curls, step-ups, front/side planks; progressive proprioception exercises; begin running straight ahead

Phase IV: 3-6 months
- WB status: full

- ROM: full

Brace: lateral padded sleeve as needed

- Exercise: Phase III + advance to open chain exercises, return to sports as tolerated