



**Amit Momaya, MD**

Orthopedic Surgeon, Chief of Sports Medicine  
Team Physician for UAB, Legion FC, and Bulls Hockey  
Highlands Appointments: 205.930.8339  
MomayaMD.com

**PATIENT NAME:**

**PROCEDURE:**

**DATE OF SURGERY:**

**PHYSICAL THERAPY ORDER – Patella Tendon Debridement and Repair**

Phase I: 0-2 weeks

- Edema and pain control
- Weight bearing status: TTWB with brace locked in extension and crutches
- ROM: 0-45 deg (ok to remove brace during ROM exercises)
- Brace: locked in extension
- Exercises: patellar mobs, calf pumps

Phase II: 2-6 weeks

- WB status: TTWB with brace locked in extension
- ROM: slowly progress to full
- Brace: locked in extension (ok to remove during ROM exercises)
- Exercises: Phase I + side lying hip/core/glutes + straight leg raises with brace locked in ext

Phase III: 6-8 weeks

- WB status: partial WB and crutches with brace unlocked
- ROM: full
- Brace: unlocked; can remove while sleeping
- Exercise: Phase II + begin calf raises, gentle stationary bike once motion allows, begin hamstring work

Phase IV: 2-4 months

- WB status: progress to WBAT without brace
- ROM: full
- Brace: wean out of brace
- Exercise: progress Phase III; avoid any squats or lunges

Phase V: 4+ months

- WB status: WBAT
- ROM: full
- Brace: none
- Exercises: cleared to progress to squats, lunges, and sports specific drills